


H

Cumulative Data Shows

Sexual contact was the most common mode of transmission, accounting for **9 out of 10 infections**.

83% of cases were among males.
For each 10 cases, 8 belonged to the 20 - 39 age group.



Of the **799 HIV positive** cases in 2012


30 were reported as AIDS cases.

29 were males


1 was female.

Philippines Data/ Fact Sheet HIV, AIDS and Sexually Transmitted Diseases

Comparison of the Distribution of Male and Female HIV cases by Age-Group and Certain Highlighted Years




Number of HIV/AIDS Cases Reported in the Philippines by the year Jan 1984 to March 2012



and at least one every three hours in **2012!**

New HIV Cases by Region as of March 2012



ROTC: Rest Of The Country

International Data

sti

448 million

new infections of
curable sexually transmitted infections
occur yearly.

trichomoniasis
chlamydia
syphilis
gonorrhoea

Some sexually transmitted infections exist without symptoms.

Sexually transmitted infections
are the main preventable cause
of infertility, particularly in
women.



70% of women

and a significant proportion of men with
gonococcal and/or chlamydial infections
experience no symptoms at all.



In pregnant women with untreated early syphilis

25%

of pregnancies result in stillbirth

14%

in neonatal death

HIV and syphilis

can also be transmitted from mother to child during pregnancy
and childbirth, and through blood products and tissue transfer.



Sources:

Philippine HIV and AIDS Registry
National Epidemiology Center, Department of
Health.

www.who.int/mediacentre/factsheets/fs110/en/index.html

http://www.who.int/features/factfiles/sexually_transmitted_diseases/facts/en/index.html

CHANGE
wellness. success. you.



Male latex condoms
**when used consistently
and correctly**

are highly effective in reducing the transmission
of HIV and other sexually transmitted infections.

HPV

One of the **most deadly sexually transmitted infections**
is the human papilloma virus.

*All cervical cancer cases are linked to genital infection
with the virus. Cancer of the cervix is the second most
common cancer in women, with about 500 000 new
cases and 250 000 deaths each year.*



The new vaccine that prevents the infection could
reduce these cervical cancer-related deaths.

CHANGE is a comprehensive workplace health and wellness campaign under the **iFLY** Keep Fit.Live Young.
Healthy Lifestyle in the Workplace Program that aims to enable young professionals to make healthy
choices to improve their well-being and overall performance.

